The Nerve! Readers Speak

Reader response: Smartphone use and primary headache: A cross-sectional hospital-based study

Joe Jacob (Thrissur, India)

Neurology: Clinical Practice December 2020 vol. 10 no. 6 e51 doi:10.1212/CPJ.00000000000997

I read the article by Uttarwar et al.¹ with interest. We are reporting special cases of bikers using mobile phone by inserting it inside their helmet over the pinnae, resulting in difficult-to-treat headache. One of our first patients was a building engineer who used to converse with his workers continuously during his ride, with phone over the ear inside the helmet. Several medications were tried for his headache until, during one of his visits, we found out his peculiar habit. Although mobile handsets are known to cause headaches, it is something inevitable in day-to-day life.^{2,3} This is because during the past few decades, several functions other than communication are integrated into it such as camera, torch light, email, internet access, videos, music, and games. Our patient reduced phone calls after our advice and also started using headsets, with which he started improving without using medications. We had several other patients with headache associated with continuous mobile phone usage.

- Uttarwar P, Vibha D, Prasad K, Srivastava AK, Pandit AK, Dwivedi SN. Smartphone use and primary headache: a cross-sectional hospital-based study. Neurol Clin Pract 2020;10:473–479.
- 2. Frey AH. Headaches from cellular telephones: are they real and what are the implications? Environ Health Perspect 1998;106:101–103.
- Wang J, Su H, Xie W, Yu S. Mobile phone use and the risk of headache: a systematic review and meta-analysis of cross-sectional studies. Sci Rep 2017;7:12595.

Copyright © 2020 American Academy of Neurology

Author response: Smartphone use and primary headache: A cross-sectional hospital-based study

Deepti Vibha (New Delhi)

Neurology: Clinical Practice December 2020 vol. 10 no. 6 e51 doi:10.1212/CPJ.00000000000998

We thank Dr. Jacob for the comment on our research article¹ and share similar observations noted in your practice. Because headache is a common neurologic problem and smartphone use is practically unavoidable in the modern era, drawing an association may be possible in your quoted case but to prescribe decreasing its use among all primary headache patients is something that requires more longitudinal studies. It may be added to the list of triggers of headache once there is more evidence. Our cross-sectional study provides the necessary association to carry this hypothesis further.

Copyright © 2020 American Academy of Neurology

Author disclosures are available upon request (ncpjournal@neurology.org)

Copyright © 2020 American Academy of Neurology. Unauthorized reproduction of this article is prohibited.

Uttarwar P, Vibha D, Prasad K, Srivastava AK, Pandit AK, Dwivedi SN. Smartphone use and primary headache: a cross-sectional hospital-based study. Neurol Clin Pract 2020;10:473–479.

Reader response: Therapeutic benefits of early electrophysiological testing in a functional neurology case

Mark Hallett, (Bethesda, MD), Kathrin LaFaver, (Chicago, IL), Carine W. Maurer, (Stony Brook, NY), Shabbir H.I. Merchant, (Charleston, SC) and Felipe Vial (Santiago, Chile) *Neurology: Clinical Practice* December 2020 vol. 10 no. 6 e52 doi:10.1212/CPJ.000000000000999

We read the article by Powell et al.¹ with interest. We advocate using clinical neurophysiology to help make the diagnosis of functional neurologic disorders (FND). Such testing can improve diagnostic certainty. We agree that the testing is valuable but for the opposite reason that the authors suggest. The clinical findings themselves were suggestive of FND. The neurophysiology points to a cerebral abnormality of motor and sensory functions. Despite the normal neuroimaging, this still could have been an episode of demyelination or another disorder. Functional disorders are often seen together with other neurologic disorders,² a situation often referred to as functional overlay. These abnormal evoked responses could be the consequence of FND. However, many more clinical neurophysiologic studies of FND have shown normal central motor and sensory conductions.³ If the findings were normal, that would be good evidence for a functional disorder. As they were abnormal, the interpretation is less certain. From a clinical point of view, the patient had positive features of FND, so it was reasonable to treat those aspects of his presentation, but not—in our view—using the clinical neurophysiology as evidence of that diagnosis.

 Stone J, Carson A, Duncan R, et al. Which neurological diseases are most likely to be associated with "symptoms unexplained by organic disease." J Neurol 2012;259:33–38.

 Hallett M. Neurophysiologic studies of functional neurologic disorders. In: Hallett M, Stone J, Carson A, eds. Functional Neurologic Disorders Handbook of Clinical Neurology, Vol 139. Amsterdam: Elsevier; 2016:61–71.

Copyright © 2020 American Academy of Neurology

Author response: Therapeutic benefits of early electrophysiological testing in a functional neurology case

Alice C. Powell, (Camperdown, New South Wales, Australia) and Michael W. Hayes (Concord, New South Wales, Australia)

We entirely agree with the comments expressed by Professor Hallett et al. in response to our article¹ regarding the use of electrophysiology to investigate probable functional neurologic disorders (FNDs). We point out in our article that electrophysiology is typically normal in FNDs and, indeed, is important evidence to support that diagnosis. However, we also mention that a small proportion of FNDs can demonstrate transient electrophysiologic abnormalities and that these seem consistent with other reported abnormal functional imaging, perfusion, and morphometric MRI studies in this group. FNDs are unlikely to be homogeneous disorders, and it may be that these exceptional cases provide insights into underlying mechanisms. This was precisely the reason for reporting this case of abnormal electrophysiology because, in our opinion, there was very robust evidence supporting a FND diagnosis: an incongruous clinical picture, pristine MRI imaging of whole spine and brain, absence of relevant signs, presence of positive phenomena, and rapid recovery.

 Powell A, Hurelbrink CB, Hayes MW. Therapeutic benefits of early electrophysiological testing in a functional neurology case. Neurol Clin Pract 2019;9:532–534.

Copyright © 2020 American Academy of Neurology

Author disclosures are available upon request (ncpjournal@neurology.org)

Powell A, Hurelbrink CB, Hayes MW. Therapeutic benefits of early electrophysiological testing in a functional neurology case. Neurol Clin Pract 2019;9:532–534.

Neurology[®]Clinical Practice

Reader response: Smartphone use and primary headache: A cross-sectional hospital-based study

Joe Jacob Neurol Clin Pract 2020;10;e51 DOI 10.1212/CPJ.0000000000000997

Updated Information & Services	including high resolution figures, can be found at: http://cp.neurology.org/content/10/6/e51.1.full.html
References	This article cites 3 articles, 1 of which you can access for free at: http://cp.neurology.org/content/10/6/e51.1.full.html##ref-list-1
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://cp.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://cp.neurology.org/misc/addir.xhtml#reprintsus

This information is current as of December 7, 2020

Neurol Clin Pract is an official journal of the American Academy of Neurology. Published continuously since 2011, it is now a bimonthly with 6 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 2163-0402. Online ISSN: 2163-0933.

