Acute Inflammatory Painful Polyradiculoneuritis

An Uncommon Presentation Related to COVID-19

Francisco Antunes Dias, MD, MSc, Ana Luiza Nunes Cunha, MD, MSc, Patrícia Maria Pedrosa Pantoja, MD, Carolina Lavigne Moreira, MD, MSc, Pedro José Tomaselli, MD, PhD, Maria Clara Zanon Zotin, MD, MSc, Francisco Antônio Colleto, MD, MSc, Soraia Ramos Cabette Fabio, MD, PhD, Octavio M. Pontes-Neto, MD, PhD, and Wilson Marques Júnior, MD, PhD

Correspondence

Dr. Dias
franciscodias.neuro@usp.br

Neurology: Clinical Practice April 2021 vol. 11 no. 2 e205-e207 doi:10.1212/CPJ.0000000000000910

Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has emerged in Wuhan, China, in December 2019, and the World Health Organization has declared it a pandemic in March 2020. Since then, severe neurologic complications associated with COVID-19 have been described, including encephalitis and stroke. Recently, it has been recognized that acute neuropathies may also occur.^{2,3} Herein, we report a patient with COVID-19 presenting an acute refractory painful polyradiculoneuritis.

PRACTICAL IMPLICATIONS

COVID-19 may be associated with neurologic complications. Patients with COVID-19 may develop an acute inflammatory painful polyradiculoneuritis that seems to respond to intravenous human immunoglobulin.

Case

A 38-year-old man presented with a 5-day history of fever, dry cough, and fatigue, followed by respiratory distress. His chest CT revealed bilateral ground-glass opacities, and he was admitted to the intensive care unit (ICU) requiring mechanical ventilation. His medical history included Class I obesity, regular alcohol, and cigarette consumption. He was treated with oseltamivir, hydroxychloroquine, azithromycin, and antibiotics for nosocomial pneumonia. Nasopharyngeal RT-PCR for SARS-CoV-2 was positive on days 1 and 12. He received a tetravalent vaccine for influenza on day 6. Sedation was discontinued 15 days after ICU admission, when he started complaining of proximally excruciating pain in both legs. Although strength and deep tendon reflexes were preserved, he was diagnosed with critical illness polyneuropathy by the ICU team, and morphine, duloxetine, and pregabalin were started. He was discharged 20 days after admission with persisting pain and negative RT-PCR.

Five days after hospital discharge, he was evaluated by a neurologist, reporting a visual analogue scale (VAS) for pain of 10 in 10 (10/10). Treatment was adjusted (duloxetine, pregabalin, amitriptyline, morphine, and vitamins supplementation), and in few days, his VAS turned 0/10. However, 1 week later, his painful manifestations returned, including tingling paresthesia and burning pain distally in lower limbs. At this time, his neurologic examination showed weakness in muscles of L5 myotomal distribution bilaterally, and pinprick-evoked pain reduced distally on both feet, accompanied by intense tactile allodynia. Deep tendon reflexes, vibration, and proprioception were normal. He had flexor plantar responses bilaterally. Cranial nerves were normal. There were no dysautonomic or sphincter abnormalities. Nerve conduction studies, including F waves, were within normal limits, and concentric needle examination showed signs of acute denervation in lumbosacral muscles innervated by right L4-L5-S1 and left L5-S1 roots (figure and table). Spinal cord MRI revealed enhancement of L5 bilaterally. CSF revealed albuminocytologic dissociation (cells: 0; protein: 76 mg/dL). An acute painful inflammatory polyradiculoneuritis was suspected, and high-dose intravenous human immunoglobulin (IVIG

MORE ONLINE

COVID-19 Resources

For the latest articles, invited commentaries, and blogs from physicians around the world

[NPub.org/COVID19](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7500000/)

Department of Neurosciences and Behavioral Sciences (FAD, ALNC, PMPP, CLM, PJT, FAC, OMP-N, WMJ), Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Department of Radiology (MCZZ), Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Department of Neurology (MCZZ), Massachusetts General Hospital, Harvard Medical School, Boston, MA; and Hospital da Unimed Ribeirão Preto (SRCF), São Paulo, Brazil.

Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the full text of this article at [Neurology.org/cp](https://www.neurology.org/cp).

(A) T1-weighted postcontrast axial with fat suppression reveals enhancement of bilateral L5 roots (black arrows). (B) T2-weighted coronal with fat suppression shows hyperintensity of bilateral L5 nerves (white arrows). (C) Chest CT with bilateral ground-glass opacities.

0.4 g/kg for 5 days) was prescribed. He has been stable (VAS 3–4/10) on the following 2 weeks. At the last visit to the neurologist, 3 weeks after IVIG infusion, the patient reported significant clinical improvement, persisting only with mild radicular pain on both legs.

Discussion

We described a case of acute inflammatory painful polyradiculoneuritis and suggest that it could be an atypical neurologic manifestation related to COVID-19 infection.

Recently, acute neuropathies have been described in relation to COVID-19. Toscano et al.⁴ reported 5 patients with COVID-19 diagnosed as having Guillain-Barré syndrome (GBS). All had severe SARS-CoV-2 infection, and neurologic symptoms emerged within the first 10 days (parainfectious profile). Four of 5 presented lower-limb flaccid weakness and areflexia that progressed

to flaccid tetraparesis/tetraplegia. Two had high CSF protein levels, and 3 had been classified as an axonal variant. Another report described a patient with severe polyradiculoneuritis leading to a locked-in syndrome.⁵ By contrast, our patient had a clinical profile restricted to lower limbs and predominantly painful, considered an acute inflammatory lumbosacral polyradiculoneuritis.

In addition, 3 Chinese patients presenting an acute painful autoimmune small-fiber neuropathy were described.⁶ They had antecedent infectious disease, presented clinically with severe pain distally in a glove-and-stockings distribution plus dysautonomia and also albuminocytologic dissociation. These key aspects (previous infectious disease, acute painful neuropathy, and albuminocytologic dissociation) are present in our patient's history and support the autoimmune nature of the peripheral nerve involvement. The symptoms evolved to the lumbosacral roots, as corroborated by MRI and EMG findings, leading us to classify it as a polyradiculoneuritis rather than a diffuse small-fiber neuropathy.

Table Nerve Conduction Studies

Nerve	CMAP							SNAP	
	L	A1	A2	A3	CV1-2	CV2-3	Fw	A	CV
R median	3.0	13.9	13.5	13.1	55.1	68.8	26.3	28.5	54.3
L median	3.1	9.2	7.9	7.5	62.0	54.8	26.1	23.4	56.3
R ulnar	2.2	7.5	6.7	6.4	64.6	65.7	26.0	12.3	61.0
L ulnar	2.6	7.8	6.9	6.3	62.3	55.9	25.8	14.8	55.3
R peroneio	4.6	4.24	4.03	3.97	41.4	44.4	48.3	7.7	40.1
L peroneio	4.4	4.67	4.21	4.19	45.8	43.9	44.3	6.7	45.0
R tibialis posterior	4.2	12.6	11.2	—	50.6	—	45.4	4.7	55.6
L tibialis posterior	3.9	16.2	10.4	—	43.6	—	44.5	5.9	59.5

Abbreviations: A = amplitude (mV for CMAP/ μ V for SAP); CMAP = compound muscle action potential; CV = conduction velocity (m/s); Fw = minimum F wave latency (ms); L = distal latency (ms); SAP = sensory action potential.

Median and ulnar nerves: 1 = wrist; 2 = elbow; 3 = axilla; Peroneal nerve: 1 = ankle; 2 = below fibula neck, 3 = above fibula neck; Tibial posterior nerve: 1 = ankle, 2 = poplitea fossa.

Neuropathic pain has already been described in patients with COVID-19.² Because no additional information was provided, it is possible that they also had a polyradiculoneuritis, similar to our case. Previous exposure to seasonal influenza vaccination should be considered a confounding causal factor, but recent epidemiologic studies question the association of this vaccine and GBS.⁷ Unfortunately, we were unable to perform RT-PCR on the CSF to exclude a direct lesion by the virus.

In conclusion, our case report suggests that patients with COVID-19 may develop an acute inflammatory painful polyradiculoneuritis that seems to respond to IVIG.

Study Funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Full disclosure form information provided by the authors is available with the full text of this article at [Neurology.org/cp](https://www.neurology.org/cp).

Publication History

Received by *Neurology: Clinical Practice* June 7, 2020. Accepted in final form July 7, 2020.

Appendix Authors

Name	Location	Contribution
Francisco Antunes Dias, MD, MSc	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Designed and conceptualized study, analyzed the data, and drafted the manuscript for intellectual content
Ana Luiza Nunes Cunha, MD, MSc	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content
Patrícia Maria Pedrosa Pantoja, MD	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content
Carolina Lavigne Moreira, MD, MSc	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content
Pedro José Tomaselli, MD, PhD	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content

Appendix (continued)

Name	Location	Contribution
Maria Clara Zanon Zotin, MD, MSc	Department of Radiology, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil; Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA	Interpreted the data and revised the manuscript for intellectual content
Francisco Antônio Colleto, MD, MSc	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Major role in the acquisition of data
Soraia Ramos Cabette Fabio, MD, PhD	Hospital da Unimed Ribeirão Preto, Ribeirão Preto, SP, Brazil	Major role in the acquisition of data
Octavio M. Pontes-Neto, MD, PhD	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content
Wilson Marques Júnior, MD, PhD	Department of Neurosciences and Behavioral Sciences, Hospital das Clínicas—Ribeirão Preto Medical School, University of São Paulo; Ribeirão Preto, SP, Brazil	Interpreted the data and revised the manuscript for intellectual content

References

- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061–1069.
- Mao L, Jin H, Wang M, et al. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol* 2020;77:1–9.
- Moriguchi T, Harii N, Goto J, et al. A first case of meningitis/encephalitis associated with SARS-coronavirus-2. *Int J Infect Dis* 2020;94:55–58.
- Toscano G, Palmerini F, Ravaglia S, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 2020;382:2574–2576.
- Pfefferkorn T, Dabitz R, von Wernitz-Keibel T, et al. Acute polyradiculoneuritis with locked-in syndrome in a patient with covid-19. *J Neurol* 2020;267:1883–1884.
- Yuki N, Chan AC, Wong AHY, et al. Acute painful autoimmune neuropathy: a variant of Guillain-Barré syndrome. *Muscle Nerve* 2018;57:320–324.
- Grave C, Boucheron P, Rudant J, et al. Seasonal influenza vaccine and Guillain-Barré syndrome: a self-controlled case series study. *Neurology* 2020;94:e2168–e2179.

Neurology® Clinical Practice

Acute Inflammatory Painful Polyradiculoneuritis: An Uncommon Presentation Related to COVID-19

Francisco Antunes Dias, Ana Luiza Nunes Cunha, Patrícia Maria Pedrosa Pantoja, et al.
Neurol Clin Pract 2021;11:e205-e207 Published Online before print July 28, 2020
DOI 10.1212/CPJ.0000000000000910

This information is current as of July 28, 2020

Updated Information & Services	including high resolution figures, can be found at: http://cp.neurology.org/content/11/2/e205.full.html
References	This article cites 7 articles, 1 of which you can access for free at: http://cp.neurology.org/content/11/2/e205.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): COVID-19 http://cp.neurology.org/cgi/collection/covid_19 EMG http://cp.neurology.org/cgi/collection/emg Guillain-Barre syndrome http://cp.neurology.org/cgi/collection/guillainbarre_syndrome Neuropathic pain http://cp.neurology.org/cgi/collection/neuropathic_pain Peripheral neuropathy http://cp.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://cp.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://cp.neurology.org/misc/addir.xhtml#reprintsus

Neurol Clin Pract is an official journal of the American Academy of Neurology. Published continuously since 2011, it is now a bimonthly with 6 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 2163-0402. Online ISSN: 2163-0933.