

Case of Guillain-Barré Syndrome After Coronavirus Disease 2019

Pablo Gómez-Porro, MD, Santiago Valenzuela-Chamorro, MD, Borja Cabal-Paz, MD, Carolina Ochoa-López, MD, Bruno García-Fleitas, MD, Edwin Ebrat-Mancilla, MD, Sabela Novo-Ponte, MD, and Raquel González-Santiago, MD

Correspondence

Dr. Gómez-Porro
gomezporrosanchez@gmail.com

Neurology: Clinical Practice December 2021 vol. 11 no. 6 e939-e941 doi:10.1212/CPJ.0000000000000917

Severe acute respiratory syndrome coronavirus (SARS-CoV-2) has spread around the world since it emerged in Wuhan, and coronavirus disease 2019 (COVID-19) has become a pandemic.¹ There is still little information regarding possible neurologic complications of this spread. Herein, we describe a patient with Guillain-Barré syndrome (GBS) preceded by COVID-19, raising the possibility of a causal relationship.

PRACTICAL IMPLICATIONS

Consider a possible association between SARS-CoV-2 infection and GBS. More epidemiologic data are required to support a causal relationship.

Case

On March 27, 2020, a 58-year-old woman with hypertension presented to our emergency department with a complaint of loss of strength in all 4 extremities. She had been diagnosed 2 months before with infiltrating breast carcinoma (cT2m pN1 cM0, Stage IIB) and received a cycle of neoadjuvant doxorubicin, cyclophosphamide, and paclitaxel, followed by right mastectomy with axillary lymphadenectomy. A second cycle of doxorubicin and cyclophosphamide was given on March 6. In addition, she had been seen by her primary care physician for complaints of dry cough, odynophagia, and dyspnea on March 2.

On March 19, she began to suffer progressive, symmetric limb weakness accompanied by paresthesia in her hands and feet. Neurologic examination disclosed bilateral facial nerve palsy, symmetric and severe tetraparesis with distal predominance, glove and stocking hypoesthesia, and global areflexia. Lung auscultation revealed bibasilar crackles. Days later, she experienced bradycardia, constipation, and urinary retention.

Blood tests on admission showed leukocytosis ($17.51 \times 10^9/L$, normal: $1.5\text{--}7.5 \times 10^9/L$), neutrophilia ($14.52 \times 10^9/L$, normal: $1.5\text{--}7.5$), thrombocytopenia ($627.00 \times 10^9/L$, normal: $150.0\text{--}400.0 \times 10^9/L$), and hyperfibrinogenemia (555.00 mg/dL , normal: $150.0\text{--}400.0 \text{ mg/dL}$). The remainder results, including lymphocytes, D-dimer, lactate dehydrogenase, and C-reactive protein, were anodyne. Serum antiganglioside antibodies were negative. CSF testing disclosed albuminocytologic dissociation (cell count $1 \times 10^6/L$, normal: $0\text{--}8 \times 10^6/L$; protein levels 114.9 mg/dL , normal: $8\text{--}43 \text{ mg/dL}$), an elevated immunoglobulin G index (0.8; normal: 0.0–0.7), negative oligoclonal bands, and negative onconeural antibodies. Brain CT showed no abnormalities. Nerve conduction studies and needle electromyography were consistent with acute demyelinating polyradiculoneuropathy with associated axonal damage (Table and Figure). The initial chest X-ray was normal, but the second showed a mixed interstitial and alveolar pattern with both lungs affected. Nasopharyngeal swabs were extracted, with a positive result for SARS-CoV-2 on reverse transcription polymerase chain reaction assay.

The patient was diagnosed with COVID-19 and GBS. She received IV immunoglobulin (IVIg, 2 g/kg) and hydroxychloroquine. Two weeks after admission, the patient

MORE ONLINE

COVID-19 Resources

For the latest articles, invited commentaries, and blogs from physicians around the world

[NPub.org/COVID19](https://www.npub.org/COVID19)

Table Motor Nerve Conduction Studies

	Distal latency, ms	Amplitude, mV	Conduction velocity, m/s	F latency, ms
Left median nerve (APB)				69.70
Wrist	12.30	2.10	—	
Elbow	25.30	1.10	18.50	
Left ulnar nerve (ADM)				57.86
Wrist	8.70	3.70	—	
Below elbow	19.35	1.60	18.80	
Above elbow	24.50	1.10	17.50	
Left peroneal nerve (EDB)				—
Ankle	19.25	0.70	—	
Fibula head	32.50	0.60	24.20	
Popliteal fossa	NR	NR	NR	
Right tibial nerve (AHB)				—
Ankle	NR	NR	NR	
Popliteal fossa	NR	NR	NR	

Abbreviations: ADM = abductor digiti minimi; AHB = abductor hallucis brevis; APB = abductor pollicis brevis; EDB = extensor digitorum brevis; NR = not recordable.

experienced significant muscle strength improvement, and her respiratory symptoms resolved.

Standard Protocol Approvals, Registrations, and Patient Consents

The authors have obtained written informed consent from the patient participating in the study.


Data Availability

The authors confirm that the data supporting the findings of this study are available within the article.

Discussion

The novel coronavirus SARS-CoV-2 is responsible for the COVID-19 pandemic. Similar to other betacoronaviruses, evidence has suggested that SARS-CoV-2 may have a neuroinvasive capacity and associated neurologic conditions. Previously, a patient with Middle East Respiratory Syndrome infection had been diagnosed with Bickerstaff encephalitis overlapping with GBS² and 4 cases of critical illness polyneuropathy/myopathy had been related to SARS-CoV.³ In April 2020, a case of possible GBS associated with SARS-

Figure Motor Nerve (Left Median and Left Ulnar) Conduction Studies



(A) Left median nerve conduction study demonstrating prolonged distal motor latency, decreased motor nerve conduction velocity, conduction block, and temporal dispersion. (B) Left ulnar nerve conduction study demonstrating prolonged distal motor latency, decreased motor nerve conduction velocity, conduction block, and temporal dispersion. (C) Increased F wave latency in the left median nerve. All of these findings are consistent with a demyelinating form of Guillain-Barré syndrome. ADM = abductor digiti minimi; APB = abductor pollicis brevis.

CoV-2 infection was reported, with a nonclassic post-infectious pattern, suggesting a possible causal relationship.⁴

Our case reports an acute demyelinating polyradiculoneuropathy in the seventeenth day of symptomatic SARS-CoV-2 infection. According to updated National Institute of Neurologic Disorders and Stroke criteria,⁵ there is a high level of certainty of a GBS diagnosis (progressive bilateral weakness of arms and legs, areflexia, autonomic dysfunction, progressive phase lasting less than 4 weeks, cranial nerve involvement, and consistent CSF and EMG findings). Moreover, as other coronaviruses have previously resulted in virus-induced neuroimmunopathology or neuropathology,⁶ the SARS-CoV-2 infection may have triggered GBS.

Given the oncological history, a possible paraneoplastic syndrome was considered as an alternative diagnosis. According to the current diagnostic criteria,⁷ it could have been a non-classic syndrome, without antibodies, within 2 years of cancer onset. Nevertheless, the existing literature suggests that these associations are possibly coincidental and specifically with breast cancer and without onconeural antibodies, only occasional. Chemotherapy drugs can also cause acute polyneuropathy, but they are typically axonal, predominantly sensitive, and without cranial nerve involvement nor dysautonomia.

In this case, respiratory and neurologic improvement was achieved after treatment with IVIg and hydroxychloroquine. COVID-19 can be severely worsened if there is an impairment of respiratory musculature. Moreover, bacterial superinfection related to dysphagia should be considered. Regarding treatment, there is already evidence that COVID-19 increases the risk of thrombosis,¹ which should be taken into account when administering IVIg. Concomitant to an active infection, plasmapheresis could be hazardous. In the matter of a novel disease, with limited scientific evidence available, clinical decisions must be individualized and previously informed consent from patients must be received.

Acknowledgment

The authors extend their gratitude to this patient and her family for their time and patience. The authors also thank Elisabeth Bassini for English language editing.

Study Funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Full disclosure form information provided by the authors is available with the full text of this article at [Neurology.org/cp](https://www.neurology.org/cp).

Publication History

Received by *Neurology: Clinical Practice* April 8, 2020. Accepted in final form July 15, 2020.

Appendix Authors

Name	Location	Contribution
Pablo Gómez-Porro, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Conceptualization of the study, acquisition of the data, analysis and interpretation of the data, and drafting of the manuscript
Santiago Valenzuela-Chamorro, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Conceptualization of the study, acquisition of the data, analysis and interpretation of the data, and drafting of the manuscript
Borja Cabal-Paz, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Conceptualization of the study, acquisition of the data, analysis and interpretation of the data, and drafting of the manuscript
Carolina Ochoa-López, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Acquisition of the data, analysis and interpretation of the data, and revision of the manuscript
Bruno García-Fleitas, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Acquisition of the data and revision of the manuscript
Edwin Ebrat-Mancilla, MD	Department of Neurophysiology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Acquisition of the data, and analysis and interpretation of the data
Sabela Novo-Ponte, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Revision of the manuscript
Raquel González-Santiago, MD	Department of Neurology, Hospital Universitario Puerta de Hierro, Majadahonda, Spain	Revision of the manuscript

References

1. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China. *N Engl J Med* 2020;382:727–733.
2. Kim JE, Heo JH, Kim HO, et al. Neurological complications during treatment of Middle East respiratory syndrome. *J Clin Neurol* 2017;13:227–233.
3. Tsai L, Hsieh S, Chao C, et al. Neuromuscular disorders in severe acute respiratory syndrome. *Arch Neurol* 2004;61:1669–1673.
4. Zhao H, Shen D, Zhou H, et al. Guillain-Barré syndrome associated with SARS-CoV-2 infection: causality or coincidence? *Lancet Neurol* 2020;19:383–384.
5. Leonhard S, Mandarakas M, Gondim F, et al. Diagnosis and management of Guillain-Barré syndrome in ten steps. *Nat Rev Neurol* 2019;15:671–683.
6. Desforges M, Le Coupance A, Dubeau P, et al. Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system? *Viruses* 2019;12:1–28.
7. Graus F, Delattre JY, Antoine JC, et al. Recommended diagnostic criteria for paraneoplastic neurological syndromes. *J Neurol Neurosurg Psychiatry* 2004;75:1135–1140.

Neurology® Clinical Practice

Case of Guillain-Barré Syndrome After Coronavirus Disease 2019

Pablo Gómez-Porro, Santiago Valenzuela-Chamorro, Borja Cabal-Paz, et al.
Neurol Clin Pract 2021;11:e939-e941 Published Online before print July 24, 2020
DOI 10.1212/CPJ.0000000000000917

This information is current as of July 24, 2020

Updated Information & Services	including high resolution figures, can be found at: http://cp.neurology.org/content/11/6/e939.full.html
References	This article cites 7 articles, 1 of which you can access for free at: http://cp.neurology.org/content/11/6/e939.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://cp.neurology.org/cgi/collection/all_clinical_neurology All clinical neurophysiology http://cp.neurology.org/cgi/collection/all_clinical_neurophysiology COVID-19 http://cp.neurology.org/cgi/collection/covid_19 Guillain-Barre syndrome http://cp.neurology.org/cgi/collection/guillainbarre_syndrome Viral infections http://cp.neurology.org/cgi/collection/viral_infections
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://cp.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://cp.neurology.org/misc/addir.xhtml#reprintsus

Neurol Clin Pract is an official journal of the American Academy of Neurology. Published continuously since 2011, it is now a bimonthly with 6 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 2163-0402. Online ISSN: 2163-0933.

